

The Management of Patients with Scabies Policy

This policy describes the processes and procedures for the management of patients with known or suspected scabies.

Key Words:	Scabies, Treatment, Infection, Prevention, Control	
Version:	7.1	
Adopted by:	Trust Policy Committee	
Date this version was adopted:	27 January 2022	
Name of Author:	Amanda Hemsley Reviewed by Claire King	
Name of Responsible Committee:	Infection Prevention and Control Group	
Please state if there is a reason for not publishing on website:	N/A	
Date issued for publication:	January 2022	
Review date:	June 2023	
Expiry date:	January 2024	
Target audience:	All LPT staff	
Type of Policy:	Clinical ✓	Non Clinical
Which Relevant CQC Fundamental Standards?		

Contents

Version control and summary of changes	3
Definitions that apply to this policy	4
1.0 Purpose of the policy	5
2.0 Summary and key points	5
3.0 Introduction	5
4.0 The management of scabies	5
4.1 Scabies	5
4.2 Classical scabies	6
4.3 Atypical scabies	7
4.4 Crusted scabies (Norwegian scabies)	7
4.5 Transmission	7
4.6 Diagnosis	8
4.7 Infection prevention and control precautions for inpatient areas	9
5.0 Specific treatment for scabies infection	9
5.1 Individual treatment	9
5.2 Recommended treatments	9
5.3 Application of treatment	10
5.4 Treatment of a patient in their own home	10
5.5 Contact tracing of an individual case	11
5.6 Further courses of treatment	11
5.7 Control of increased incidences and outbreaks	11
5.8 Movement of symptomatic patients from an affected unit	12
5.9 Health care personnel	12
6.0 Training	13
7.0 References and bibliography	13
Appendix 1: Contact details	14
Appendix 2: Management of an Outbreak of Scabies flowchart..	15
Appendix 3: Privacy Impact Assessment	16
Appendix 4: Stakeholders and Consultation.....	17
Appendix 5: NHS Consultation.....	18
Appendix 6: Due Regard Screening.....	19

Version Control and Summary of Changes

Version	Date	Comment
1.	June 2010	Review of current interim guidelines. Circulated for comments to all members of LCCHS Infection Control Sub Committee and Infection Control Link Staff Adults and Children's Community Health Service
2.	July 2010	Amendments following consultation process. Revisions to incorporate requirements of NHSLA Standards
3.	July 2010	Re-circulated for comments Amendments following consultation process. Forwarded to LCCHS Clinical Governance Committee for approval.
4.	August 2011	Harmonised in line with LCRCHS, LPT, LCCHS (Historical organisations)
5.	June 2015	Review and update in line with organisation policy requirements
6.	May 2018	Review and update in line with organisation policy requirements
7.	October 2021	Review and update in line with organisation policy requirements
7.1	October 2023	Change made to occupational health number

For further information contact: Infection Prevention and Control Team
01162951668

Definitions that apply to this Policy

Allergy	A condition of increased sensitivity to a substance (an allergen) considered harmless to most people.
Consultant in Public Health	A consultant who is knowledgeable in Infectious Diseases
Contact Tracing	The identification and diagnosis of persons who may have come into contact with an infected person.
Hyperkeratosis	Is a thickening of the outer layer of the skin, which contains a tough protein called keratina. This thickening is often part of the skin's normal protection against rubbing, pressure and other forms of local irritation and causes calluses and corns on the soles of the feet or of the hands.
Infection	An organism presents at a site and causes an inflammatory response or where an organism is present in a normally sterile site.
Isolation	When a patient is cared for in a separate area or room due to them having an infection that may be detrimental to other individual's health. Or when the patient may be vulnerable to infection.
Outbreak	The occurrence of two or more cases of the same infection linked in time and place or, the situation when the observed number of cases exceeds the number expected.
Personal Protective Equipment (PPE)	Specialised clothing or equipment worn by employees for protection against health and safety hazards. Gloves, aprons, gowns, masks and eye protection
Symptomatic	Physical or mental sign of the disease.
Treatment	Care provided to improve a situation (especially medical procedures or applications that are intended to relieve illness or injury)

1.0 Purpose of the policy

The purpose of this policy is to inform all healthcare staff within Leicestershire Partnership Trust (LPT) who are involved in the care of patients that develop or suffer from symptoms or infection of scabies, the process and management of the infection. When individuals are in close and frequent contact with each other, infectious diseases can spread rapidly both within a healthcare setting and in the community.

Staff working within LPT provide a number of health services to the wider community. This document provides information on the processes required when treating, managing or giving advice regarding scabies. It supports the prevention of cross infestation amongst the wider population. This policy has been produced in accordance with published evidence and national best practice guidelines. As a duty of care LPT must ensure that staff are given guidance as to the appropriate steps they need to undertake to ensure that they can protect the patients within their care.

2.0 Summary and key points

The provision of healthcare carries with it inherent risks to the health care worker. The purpose of this policy is to ensure that all staff are aware of their responsibilities for safe practice in relation to the management of scabies and take the appropriate precautionary measures to protect themselves, their co-workers, and their patients.

3.0 Introduction

This policy has been developed to provide organisational wide guidance for the management and treatment of cases of scabies. It defines scabies and discusses presentation, transmission, and diagnosis. Infection prevention control precautions are then highlighted with particular reference to the management and treatment of scabies with the overall aim being to reduce the risk of transmission.

Appendices detail contacts for advice and a flowchart for the management of an increased incidence/outbreak of scabies have been included for reference.


4.0 The management of scabies

4.1 Scabies

Scabies is an infestation caused by a mite known as *Sarcoptes scabiei*, which burrows into the skin and lays its eggs. An allergy to mite eggs and the faecal droppings produced is responsible for symptoms of intense itching and a characteristic rash.

Scabies is known to cause sporadic cases and outbreaks in hospitals, nursing/residential homes, schools and any other communal care environments where people have direct prolonged skin contact. Effective control is dependent on early diagnosis, adequate treatment of cases and contacts and the prevention of further spread

The Scabies mite is:



- Oval in shape and measures 0.2 – 0.4 mm in length
- The body is covered with fine lines and long hairs
- Is blind
- Has 8 legs when an adult
- Usually lives for 30 – 60 days
- Cannot jump
- Females burrow into the skin to lay eggs
- Males live on the surface of the skin
- Life cycle of 4 – 6 weeks
- Not related to hygiene standards

Scabies is most common:

- In females
- In people aged 10 – 19 years
- In areas with high levels of social deprivation
- During winter

The appearance and severity of symptoms are strongly influenced by the immune status of the affected person. Scabies may present as follows:

4.2 Classical scabies

This is the form of scabies generally found in healthy people with a normal immune system.

- The number of mites present in classical scabies is small (15-20) and spread is usually by direct physical contact.
- Burrows appear as irregular, raised discolored lines 0.5 to 1.5cm long in the skin, which are not always visible to the naked eye.
- Often a bilateral symmetrical rash can be seen in areas such as the midriff, inner thighs and axillae.
- There is extreme itching, especially at night.
- The sites of the rash and burrows do not necessarily correspond



4.3 Atypical scabies

Atypical scabies occurs in any person with immature or impaired immune response. Many mites may be present in atypical scabies and symptoms may be variable. Scaling or crusting of the skin may be present but is usually slight. Itching may also be very slight or even absent

A high proportion of atypical cases occur in the elderly, and particularly within residential/nursing homes and elderly units.

4.4 Crusted scabies (Norwegian scabies)

This form of scabies is extremely infectious and occurs in those whose immune systems are severely impaired.

- Hyperkeratotic skin lesions appear as hardened crusts containing thousands or millions of mites.
- Itching may be slight or absent.
- Skin becomes crusted especially on palms, soles, nail beds, wrists, buttocks and penis. The whole body can be affected including the head and scalp.



4.5 Transmission

Scabies is mainly spread from person to person by direct, prolonged skin contact with an infected person, such as prolonged holding of hands

Scabies mites rapidly die, once away from the human body and therefore clothing and linen etc. is not the main route for transmission. However, mites shed in skin scales, can live in the environment longer and therefore the risk of spread through contact with soft furnishings / carpets clothing and linen is increased with heavy skin shedders. Pets do not spread scabies.

The length of time between contact with an affected person and developing signs of scabies (i.e. itching and a rash) is between four and six weeks, if this is the first infection. For re-infection symptoms appear within 48 hours.

4.6 Diagnosis

Diagnosis is by identification of the mite, eggs, or faecal matter from skin scrapings. A clinical diagnosis may be made by a combination of severe itching especially at night and a typical or atypical distribution of a rash in persons who have had skin to skin contact with suspected or diagnosed cases.

Another hallmark of scabies is the appearance of track-like burrows in the skin. These raised lines are usually grayish-white or skin-colored. They are created when female mites tunnel just under the surface of the skin. After creating a burrow, each female lays 10 to 25 eggs inside.



If clinical diagnosis is difficult, obtaining skin scrapings for the scabies mites or their faecal pellets may be helpful. Skin scales may be sent by the clinician caring for the patient in a closed, sealed container (e.g. universal container) to the Microbiology Laboratory University Hospitals Leicester NHS Trust for microscopy for scabies.

Once the diagnosis is confirmed a dermatology opinion may be necessary.

4.7 Infection prevention and control precautions for inpatient areas

All patients in communal health care environments who are suspected of having or have been diagnosed as having scabies should be cared for in source isolation until treatment has been administered and washed off, or an alternative diagnosis has been made. The infection prevention and control team must be informed of the patient as soon as possible. Two courses of treatment are required and must be administered a week apart.

Precautions must continue until the two courses of treatment have been administered and washed off or an alternative diagnosis made.

Disposable nitrile gloves and plastic aprons must be worn when in contact with the patient, bed linen and patients clothing. The use of personal protective equipment (PPE) must be used until the second course of treatment has been administered and washed off.

Hand washing with liquid soap and water must be carried out after contact with the patient and their environment. Hands must be dried thoroughly with single use disposable paper towels.

Linen should be treated as infected until the patient has had the second course of treatment. Clean clothing and bed linen should be available after treatment has been washed off.

4.7. 1 Environmental cleaning

The environment and equipment must be cleaned with chlor-clean as per the Cleaning and Decontamination policy.

Good environmental cleaning is an integral and important component preventing healthcare-associated infections within inpatient areas, other healthcare settings and non-health care buildings where health care is delivered. The environment must be visibly clean, free from dust and soil, with the overall appearance being acceptable to patients, visitors, and staff. A clean environment reflects the quality of care, structure, and efficient function of Leicestershire Partnership NHS Trust (LPT).

The environment is known to play an important role in cross infection which can lead to increased incidents or outbreaks of infection. Door handles, flush handles, taps, toilet roll holders etc. have all been implicated at some point in this potential risk. Therefore, accumulation of dust, dirt and liquid residues may increase the risks and must be reduced to the minimum. This can be achieved by regular cleaning and by good design features in buildings, fittings, and fixtures. A program for monitoring the standard of cleaning is in place across all clinical settings via the environmental audits and cleaning audit program.

5.0 Specific treatment for scabies infection

5.1 Individual treatment

Individual treatment should be given if scabies is diagnosed.

5.2 Recommended treatments

Permethrin 5% (Lyclear Dermal Cream), - 30g (low toxicity, non-irritant). Leave for 8-12 hours.

Malathion 0.5% (Derbac-M) – leave for 24 hours

Malathion 0.5% is the treatment of choice in pregnancy, during breast feeding and for infants under 2 months.

Permethrin 5% is the treatment of choice in children over the age of 2 months and under the age of 6 months

For each treatment the following amounts will be required:

- 200ml of lotion or 30g cream, for an average sized person.
- More than 200ml of lotion or more than 30g of cream may be required for a larger person or for a patient where there is a need for frequent washing

5.3 Application of treatment

The lotion or cream should be applied to all skin surfaces starting at the jawline and around behind the ears, extending to the soles of the feet.

It should be applied to cool dry skin and never after a hot bath. The lotion or cream should be reapplied to skin whenever it has been washed during the treatment time (as these areas will not undergo the recommended time for treatment and may promote reinfection). Lotion or cream used should then be allowed to dry before the person dresses.

The head and scalp may be affected and should be carefully examined for any signs of scabies and be treated if present after seeking medical advice.

Following the recommended time, depending on the type of medication being used the lotion or cream should be **washed off thoroughly** with plenty of water. This should be done preferably by a shower (or bath if a shower is not an option).

Itching may persist and the rash may be present for up to 2 weeks after treatment. The use of calamine lotion, crotamiton (Eurax) or antihistamines may be helpful in the management of itching.

For inpatient facilities, clothes, underwear and bedding used in the 48 hours prior to and during the treatment should be placed into a **red soluble bag** and securely tied before being placed into a **white outer plastic bag** (Please refer to Infection prevention and Control Policy for the Management of Linen and Laundry).

Within community facilities and/or primary care clothes, underwear and bedding used in the 48 hours prior to and during the treatment should be laundered in a hot wash (over 50°C). If this is not possible, the laundry should be stored in a sealed plastic bag for 72 hours prior to being washed, when it can then be washed following the washing instructions on the garment. This laundry must not come into contact or be stored with clean clothing or linen.

Floors and upholstery should be vacuumed after the removal of contaminated linen. Clean clothing and bedding should be used after the lotion has been washed off.

If hyperkeratotic lesions are present, then specialist advice on appropriate management is needed from a dermatologist.

Inform the Infection Prevention and Control Team. Please refer to Contacts for advice (Appendix 1) of all suspected and diagnosed cases of scabies.

5.4 Treatment of a patient in their own home

An initial assessment should be made by the clinician caring for the patient as to whether the client is suitable for treatment in their own home.

The patient should be mobile/flexible enough to be able to have the treatment applied to the whole of their body and be able to have the treatment washed off preferably in a shower or bath. However, where a bath or shower is not available or possible, then lotion should be thoroughly washed and rinsed off with care. For further advice on the management of individual patients please contact the Infection Prevention and Control

Team.

Where patients already attend social services day centers for bathing, the treatment should be applied by the appropriate health care staff prior to them attending the care home.

Arrangements may be made in advance with the Care Home Staff for the treatment to be washed off. Patients should not attend day centers until the treatment has been completed.

In the case of children attending school or nursery the child must be kept away from the school or nursery until treatment has been completed.

For patients who are difficult to treat at home, advice should be sought from the Infection Prevention and Control Team.

5.5 Contact tracing of an individual case

Following consultation with the Medical Practitioner, Consultant in Public Health or Infection Prevention and Control Team the treatment of close contacts should be arranged by the individual who has been diagnosed with scabies. A risk assessment will be made to ascertain the patient's current status and the level of contact others have with the patient.

In the case of children, contacts should have one treatment at the same time as the second treatment of the affected case. Treatment should include the entire household and close contacts.

Close contacts may include partners, parents, siblings, young dependent children, carers and staff caring for the affected patient.

5.6 Further courses of treatment

- Further treatment may be necessary depending on the extent and severity of the scabies infection.
- In the management of severely infected patients at least 3 treatments will be necessary.
- If multiple treatments are prescribed for an individual, they should be at least 7 days apart.
- Where treatment has failed, advice should be sought from the Infection Prevention and Control Team.
- Persons are classed as non-infectious when their treatment regimen has finished, and the medication washed off.

5.7 Control of increased incidences and outbreaks (also see Appendix 2 – increased incident/outbreak flowchart)

If two or more cases of scabies are detected in the same inpatient facility and are linked in time, (2 weeks or less apart) then the Infection Prevention and Control Team must be notified (appendix 2).

If two or more cases of scabies are detected in the same nursing/residential home,

closely linked in time (2 weeks or less apart), then the Infection Prevention and Control team for the local authority and Public Health England, East Midlands Health Protection Team must be notified.

If two or more cases of scabies are detected in the same school/nursery, closely linked in time (2 weeks or less apart), then the Infection Prevention and Control Team and Public Health England, East Midlands Health Protection Team must be notified.

If an increased incident or outbreak is suspected, referral to a dermatologist is required to urgently confirm diagnosis of scabies. The referral for patients should be made by their clinician i.e. medical practitioner, ANP. A referral for staff should be made by an Occupational Health Doctor (Please refer to Contacts for Advice, Appendix 1). Once an increased incident/outbreak has been identified the increased incident/outbreak policy will be implemented.

Planned coordinated treatment is essential and where possible, individuals should be treated at the same time to prevent the likelihood of re-infection.

Where staff require treatment, this will be managed by the Occupational Health Department.

In conjunction with the Dermatologist, the Consultant in Public Health (please refer to Contacts for Advice, Appendix 1) will decide who needs treatment and the treatment regimen to be carried out, taking into account the following information: -

- a) The number of symptomatic patients in the affected unit.
- b) The number of symptomatic staff working in the unit.
- c) The total number of patients and staff within the unit with or without symptoms.
- d) The severity of symptoms of each affected individual.

From this information the Consultant in Communicable Disease will then decide on whether to treat symptomatic individuals only or all patients and all staff based in the unit.

Close contacts must be treated at the same time as the last treatment of the symptomatic individual.

5.8 Movement of symptomatic patients from an affected unit

Symptomatic patients from an affected unit should ideally not be transferred or discharged to other communal health care environments including other hospital units, residential or nursing homes until coordinated treatment has been given and washed off. In circumstances where patient movement is necessary communication is vital between the two areas so that appropriate infection prevention and control precautions can be adopted. In such instances the Essential Steps Inter-Hospital Transfer Form is to be used (Please refer to appendix 3).

5.9 Health care personnel

It is the responsibility of the Occupational Health Department to manage the treatment of staff when there is a case of occupationally acquired scabies.

A referral to a dermatologist may be necessary to confirm the diagnosis of scabies in staff. In this case, the Occupational Health Department managing the healthcare personnel should contact the dermatology department for urgent assistance in confirming diagnosis.

The Occupational Health Department will prescribe treatment for staff and close contacts of staff as necessary. LPT will meet the cost of the necessary treatment that has been prescribed by the Occupational Health Department.

6.0 Training

There is no training requirement identified within this policy

7.0 References and bibliography

The supporting Infection Prevention and control policies can be located at:

<http://www.leicspart.nhs.uk/SupportServices>

[Management of Linen and Laundry in Community Health Services, Inpatient Services and Primary Care](#)

[Management of a Patient requiring source isolation in Community Health Services, Inpatient Facilities and Primary Care](#)

[Personal Protective Equipment for use in Healthcare Policy](#)

Hand hygiene Policy (including Bare Below the Elbows)

British National Formulary September 2020

National Institute for Health and Care Excellence (NICE) - Clinical skill Summaries: Scabies, <https://cks.nice.org.uk/scabies#!topicsummary>

Scabies – NHS.UK <http://www.nhs.uk/conditions/Scabies>

Public Health England www.phe.gov.uk
(Listed under S for scabies)


Wilson, J (2006) Infection Control in Clinical Practice. London: Bailliere Tindall.

Appendix 1

Contact details		
Name	Address	Telephone number
Public Health England East Midlands Health Protection Team	<p>East Midlands Health Protection Unit Seaton House City Link Nottingham NG2 4LA</p> <p>Email: emhpt@phe.gov.uk</p> <p>Area's covered: Leicestershire, Rutland. Lincolnshire, Nottinghamshire, Derbyshire</p>	<p>☎0344 225 4524 (option 1)</p> <p>Fax: 0116 263 0</p>
Occupational Health Department	Baldwin Lodge Glenfield General Hospital	☎0116 2585307
Infection Prevention and Control Team Leicestershire Partnership Trust	Loughborough hospital Epinal way Loughborough LE11 5JY	☎0116 2951668

Appendix 2

Management of an increased incident/outbreak of scabies



Appendix 3

PRIVACY IMPACT ASSESSMENT SCREENING

<p>Privacy impact assessment (PIAs) are a tool which can help organisations identify the most effective way to comply with their data protection obligations and meet individual's expectations of privacy. The first step in the PIA process is identifying the need for an assessment.</p> <p>The following screening questions will help decide whether a PIA is necessary. Answering 'yes' to any of these questions is an indication that a PIA would be a useful exercise and requires senior management support, at this stage the Head of Data Privacy must be involved.</p>			
Name of Document:	The management of patients with scabies policy		
Completed by:	Claire king		
Job title	Infection Prevention and Control Nurse	Date	20/07/2021
			Yes / No
1. Will the process described in the document involve the collection of new information about individuals? This is information in excess of what is required to carry out the process described within the document.			No
2. Will the process described in the document compel individuals to provide information about themselves? This is information in excess of what is required to carry out the process described within the document.			No
3. Will information about individuals be disclosed to organisations or people who have not previously had routine access to the information as part of the process described in this document?			No
4. Are you using information about individuals for a purpose it is not currently used for, or in a way it is not currently used?			No
5. Does the process outlined in this document involve the use of new technology which might be perceived as being privacy intrusive? For example, the use of biometrics.			No
6. Will the process outlined in this document result in decisions being made or action taken against individuals in ways which can have a significant impact on them?			No
7. As part of the process outlined in this document, is the information about individuals of a kind particularly likely to raise privacy concerns or expectations? For examples, health records, criminal records or other information that people would consider to be particularly private.			No
8. Will the process require you to contact individuals in ways which they may find intrusive?			No
<p>If the answer to any of these questions is 'Yes' please contact the Head of Data Privacy Tel: 0116 2950997 Mobile: 07825 947786 Lpt-dataprivacy@leicspart.secure.nhs.uk In this case, ratification of a procedural document will not take place until approved by the Head of Data Privacy.</p>			
IG Manager approval name:			
Date of approval			

Acknowledgement: Princess Alexandra Hospital NHS Trust

Appendix 4

Key individuals involved in developing the document

	Designation
Amanda Hemsley	Lead Infection Prevention and Control Nurse
Antonia Garfoot, Laura Brown Andy Knock Clarissa Swann Claire King	Infection Prevention and Control Team

Circulated to the following individuals for consultation

Name	Designation
Anne Scott	Executive director of nursing, AHPS and quality
Emma Wallis	Associate director of nursing & professional practice
Claire Armitage	Lead nurse for community AMH
Alison O'Donnell	Interim head of learning and development
Michelle Churchard	Head of nursing AMH/LD services
Louise Evans	Deputy head of nursing FYPC/LD services
Kam Palin	Occupational health nurse
Tejas Khatau	Lead pharmacist FYPC
Jane Martin	Acting deputy head nursing DMH
Katie Willetts	Senior Nurse, specialist nursing FYPC
Bernadette Keavney	Head of trust health and safety compliance
Maureen Poyzer	Health and safety advisor
Cheryl Shuttleworth	Facilities manager
Helen Walton	Estates and facilities property manager
Clare Pope	LD Modern matron Bradgate unit
Sarah Latham	Deputy head of nursing community hospitals
Elizabeth Compton	Senior matron AMH Bradgate unit

The NHS Constitution

The NHS will provide a universal service for all based on clinical need, not ability to pay. The NHS will provide a comprehensive range of services

Shape its services around the needs and preferences of individual patients, their families and their carers	<input type="checkbox"/>
Respond to different needs of different sectors of the population	<input type="checkbox"/>
Work continuously to improve quality services and to minimise errors	<input type="checkbox"/>
Support and value its staff	<input type="checkbox"/>
Work together with others to ensure a seamless service for patients	<input type="checkbox"/>
Help keep people healthy and work to reduce health inequalities	<input type="checkbox"/>
Respect the confidentiality of individual patients and provide open access to information about services, treatment and performance	<input type="checkbox"/>

Due Regard Screening Template

Section 1	
Name of activity/proposal	Infection Prevention and Control Overarching Policy
Date Screening commenced	17 May 2018
Directorate / Service carrying out the assessment	Enabling. Infection Prevention and Control Team
Name and role of person undertaking this Due Regard (Equality Analysis)	Amanda Hemsley, Lead Infection Prevention and Control Nurse
Give an overview of the aims, objectives and purpose of the proposal:	
AIMS:	
To provide clear guidance to Trust staff on their responsibilities in relation to infection prevention and control.	
OBJECTIVES:	
This policy clearly identifies the aims and goals for infection prevention and control within Leicestershire Partnership Trust, thereby providing a coherent strategic objective. This policy should be reviewed whenever there is a need to adapt to the changing regulatory environment or in response to ongoing risk assessment to ensure a safe environment exists for all patients, visitors and staff.	
Section 2	
Protected Characteristic	If the proposal/s have a positive or negative impact please give brief details
Age	This document provides guidance on the roles and responsibilities of all staff working within the trust in relation to the prevention and control of infection. Therefore the correct implementation of this policy will help reduce any adverse effect irrespective of any protected characteristic and is therefore equality neutral
Disability	
Gender reassignment	
Marriage & Civil Partnership	
Pregnancy & Maternity	
Race	
Religion and Belief	
Sex	
Sexual Orientation	
Other equality groups?	

Section 3


Does this activity propose major changes in terms of scale or significance for LPT? For example, is there a clear indication that, although the proposal is minor it is likely to have a major affect for people from an equality group/s? Please tick appropriate box below.

Yes		No	
High risk: Complete a full EIA starting click here to proceed to Part B		Low risk: Go to Section 4.	✓

Section 4

If this proposal is low risk please give evidence or justification for how you reached this decision:

This policy is the overarching policy for all subsequent infection prevention and control policies. The policies take into consideration the needs of patients and staff and the safeguarding of same. It follows government legislation and relevant bodies have been consulted prior to the development of any policies prior to having them agreed at trust board level.

Signed by reviewer/assessor		Date	3 November 2020
------------------------------------	--	-------------	-----------------

Sign off that this proposal is low risk and does not require a full Equality Analysis

Head of Service Signed		Date	
-------------------------------	--	-------------	--